

Results on Convergence in Norm of Exponential Product Formulas and Pointwise of the Corresponding Integral Kernels

Takashi Ichinose and Hideo Tamura

Abstract. For the last one and a half decades it has been known that the exponential product formula holds also *in norm* in nontrivial cases. In this note, we review the results on its convergence in norm as well as pointwise of the integral kernels in the case for Schrödinger operators, with error bounds. Optimality of the error bounds is elaborated.

Mathematics Subject Classification (2000). Primary 47D08; Secondary 47D06, 41A80, 81Q10, 35J10.

Keywords. Exponential product formula, Trotter product formula, Lie–Trotter product formula, Trotter–Kato product formula, pointwise integral kernel convergence, evolution group and semigroup, Schrödinger operator.

1. Introduction

The Trotter product formula, Trotter–Kato product formula or exponential product formula is usually a product formula which in strong operator topology approximates the group/semigroup with generator being a sum of two operators. It is often a useful tool to study Schrödinger evolution groups/semigroups in quantum mechanics and to study Gibbs semigroups in statistical mechanics.

To think of a typical case, let A and B be selfadjoint operators in a Hilbert space $\mathcal{H}$ with domains $D[A]$ and $D[B]$ and $H := A + B$ their operator sum with domain $D[H] = D[A] \cap D[B]$. Assume that H is selfadjoint or essentially selfadjoint on $D[H]$ and denote its closure by the same H . Then Trotter [44] proved the unitary product formula

$$\begin{aligned} [e^{-itB/2n} e^{-itA/n} e^{-itB/2n}]^n - e^{-itH} &\rightarrow 0, \quad \text{strongly,} \\ [e^{-itA/n} e^{-itB/n}]^n - e^{-itH} &\rightarrow 0, \quad \text{strongly,} \quad n \rightarrow \infty, \end{aligned}$$

and also, when A and B are nonnegative, the selfadjoint product formula

$$\begin{aligned} [e^{-tB/2n} e^{-tA/n} e^{-tB/2n}]^n - e^{-tH} &\rightarrow 0, \quad \text{strongly,} \\ [e^{-tA/n} e^{-tB/n}]^n - e^{-tH} &\rightarrow 0, \quad \text{strongly.} \quad n \rightarrow \infty, \end{aligned}$$

The convergence is *locally uniform*, i.e. uniform on compact t -intervals, respectively in the real line $\mathbf{R}$ and in the closed half line $[0, \infty)$. Kato [29] discovered the latter selfadjoint product formula to hold also for the form sum $H := A \dot{+} B$ with form domain $D[H^{1/2}] = D[A^{1/2}] \cap D[B^{1/2}]$, which we assume for simplicity is dense in $\mathcal{H}$. However, it remains to be an open problem whether the unitary product formula for the form sum holds.

However, since around 1993 we have begun to know that selfadjoint product formulas converge even in (*operator*) *norm*, though in some special cases, by the following two first results. Rogava [37] proved, when B is A -bounded and $H = A + B$ is selfadjoint, among others, the abstract product formula that

$$\| [e^{-tA/n} e^{-tB/n}]^n - e^{-tH} \| = O(n^{-1/2} \log n), \quad n \rightarrow \infty,$$

locally uniformly in $[0, \infty)$. Helffer [13] proved, when $H := -\Delta + V(x)$ is a Schrödinger operator in $L^2(\mathbf{R}^d)$ with nonnegative potential $V(x)$ satisfying $|\partial_x^\alpha V(x)| \leq C_\alpha$ ($|\alpha| \geq 2$) so that H is selfadjoint on the domain $D[-\Delta] \cap D[V]$, the symmetric product formula that

$$\| [e^{-tV/2n} e^{-t(-\Delta)/n} e^{-tV/2n}]^n - e^{-tH} \| = O(n^{-1}), \quad n \rightarrow \infty,$$

locally uniformly in $[0, \infty)$. Many works were done to extend these results before 2000, e.g. in [5, 20, 22, 32, 33, 35] for the abstract product formula, [9, 10, 17, 18, 19, 41] for the Schrödinger operators, and after that, e.g. in [23, 27, 16], [3, 4, 6, 7] for the abstract product formula. In most of them, use was made of operator-theoretic methods, though of a probabilistic method in [17, 18, 19, 41].

In this note, we want to describe more recent results on convergence in norm for exponential product formulas and also pointwise of the corresponding integral kernels, mainly based on our works since around 2000, [23, 27, 24, 25, 26]. As for the error bounds, although it is easy to see by the Baker–Campbell–Hausdorff formula (e.g. [45], [40]) that with both operators A and B being bounded, the *nonsymmetric* product formula has an optimal error bound $O(n^{-1})$ while the *symmetric* one does $O(n^{-2})$, it was shown in [27] that even the *symmetric* product formula has an optimal error bound $O(n^{-1})$ in general, if both A and B are unbounded. However, in [25] (cf. [26]), a better upper sharp error bound $O(n^{-2})$ has been obtained for the *symmetric* product formula with the Schrödinger operator $-\Delta + V(x)$ having nonnegative potentials $V(x)$ growing polynomially at infinity, in spite that both $-\Delta$ and V are unbounded operators. In this note we mention, with a sketch of proof, a latest complementary result [2] which settles the sharp optimal error bound is in fact $O(n^{-2})$ with the symmetric product formula for the harmonic oscillator, by estimating the error not only from above but also from below, in norm as well as pointwise.

Theorems are described in Section 2. Optimality of error bounds is discussed separately in Section 3. The idea of proof is briefly mentioned in Section 4. In Section 5 we give concluding remarks, and also refer to a connection of the exponential product formula with the Feynman path integral.

It should be also noted that in almost the same context with the notion of norm ideals (e.g. [12], [38]) we are able to deal with the trace norm convergence as in [46, 30, 31, 34, 14, 21, 42]. For an extensive literature on this we refer to [47].

The content of this note is an expanded version of the lecture entitled “On convergence pointwise of integral kernels and in norm for exponential product formulas” given by T.I. at the International Conference “*Modern Analysis and Applications* (MAA 2007)”, Odessa, Ukraine, April 9–14, 2007, which is a slightly extended version of the lecture (unpublished) given at the Conference on “*Heat Kernel in Mathematical Physics*”, Blaubeuren, Germany, November 28–December 2, 2006.

2. Theorems

We begin with our result which extends ultimately Rogava and Helffer’s.

Theorem 2.1. (*Ichinose-Tamura-Tamura-Zagrebnov 2001*[23, 27]) *Let A and B be nonnegative selfadjoint operators, and assume $H = A + B$ is selfadjoint on $D[H] = D[A] \cap D[B]$. Then as $n \rightarrow \infty$,*

$$\| [e^{-tB/2n} e^{-tA/n} e^{-tB/2n}]^n - e^{-tH} \| = O(n^{-1}), \quad (2.1)$$

$$\| [e^{-tA/n} e^{-tB/n}]^n - e^{-tH} \| = O(n^{-1}). \quad (2.2)$$

The convergence is locally uniform in the closed half line $[0, \infty)$, while on the whole half line $[0, \infty)$, if H is strictly positive, i.e. $H \geq \eta I$ for some $\eta > 0$. The error bound $O(n^{-1})$ in (2.1) and (2.2) is optimal.

We can go beyond this result. First, focussing on the Schrödinger operator $-\Delta + V(x)$, we ask whether norm convergence implies pointwise convergence of integral kernels. The answer is yes, though strong convergence does not. This problem is discussed for Schrödinger operators with potentials of polynomial growth (Theorem 2.2), with positive Coulomb potential (Theorem 2.3), and also for the Dirichlet Laplacian (Theorem 2.4). Pointwise convergence of integral kernels for Schrödinger semigroups is important, because it gives a time-sliced approximation to the imaginary-time Feynman path integral.

Next, we ask, for the unitary exponential/Trotter product formula, whether there are nontrivial cases where it converges in norm, though it does not in general hold (see [15]). The answer is yes. In fact, it holds for the Dirac operator and relativistic Schrödinger operator (Theorem 2.5).

Let $H = H_0 + V := -\Delta + V(x)$ with $V(x)$ a real-valued function. By $K^{(n)}(t, x, y)$ we denote the integral kernel of $[e^{-tH_0/2n} e^{-tV/n} e^{-tH_0/2n}]^n$, and by $e^{-tH}(x, y)$ that of e^{-tH} .

Theorem 2.2. (*Ichinose-Tamura 2004 [25]*) (*positive potential of polynomial growth*) Assume that $V(x)$ is in $C^\infty(\mathbf{R}^d)$, bounded below and satisfies $|\partial_x^\alpha V(x)| \leq C_\alpha \langle x \rangle^{m-\delta|\alpha|}$ with some $0 < \delta \leq 1$ ($\langle x \rangle = (1 + x^2)^{1/2}$).

(i) (*In norm*)

$$\| [e^{-tH_0/2n} e^{-tV/n} e^{-tH_0/2n}]^n - e^{-tH} \|_{L^2} = O(n^{-2}), \quad (2.3)$$

locally uniformly in the open half line $(0, \infty)$.

(ii) (*Integral kernel*)

$$\begin{aligned} [K^{(n)}(t, x, y) - e^{-tH}(x, y)] &= O(n^{-2}), \\ \text{in } C^\infty(\mathbf{R}^d \times \mathbf{R}^d)\text{-topology, locally uniformly in } (0, \infty), \end{aligned} \quad (2.4)$$

i.e. together with all x, y -derivatives.

This theorem improves the result of Takanobu [41], who used a probabilistic method with the Feynman–Kac formula (see Sect. 5) to show uniform pointwise convergence of the integral kernels, roughly speaking, with error bound $O(n^{-\rho/2})$, if $V(x)$ satisfies $V(x) \geq C(1 + |x|^2)^{\rho/2}$ and $|\partial_x^\alpha V(x)| \leq C_\alpha(1 + |x|^2)^{(\rho-\delta|\alpha|)_+/2}$ for some constants $C, C_\alpha \geq 0$ and $\rho \geq 0, 0 < \delta \leq 1$. The claim of Theorem 2.2 is a little bit sharpened in Theorems 3.1 and 3.2, in the next section, in the case of the harmonic oscillator.

Theorem 2.3. (*Ichinose-Tamura 2006 [26]*) (*positive Coulomb potential*) Let $H := -\Delta + V(x)$ with $V(x) \geq 0$. Assume that $V(-\Delta + 1)^{-\alpha}: L^2(\mathbf{R}^d) \rightarrow L^2(\mathbf{R}^d)$ is bounded for some $0 < \alpha < 1$, and that $V \in C^\infty$ near a neighbourhood U of both p and q (after $p, q \in \mathbf{R}^d$ taken). Then

$$\begin{aligned} [K(t/n)^n(x, y) - e^{-tH}(x, y)] &= O(n^{-1}), \\ \text{in } C^\infty(U)\text{-topology, locally uniformly in } (0, \infty). \end{aligned} \quad (2.5)$$

The condition is satisfied if V is in $L^2(\mathbf{R}^3) + L^\infty(\mathbf{R}^3)$, in particular, if V is the positive Coulomb potential $1/|x|$. We don't know what happens at the singularities of $V(x)$.

Theorem 2.4. (*Ichinose-Tamura 2006 [26]*) (*Dirichlet Laplacian*) Let $\Omega \subset \mathbf{R}^d$ be a bounded domain with smooth boundary and χ_Ω the indicator function of Ω . Let $H_0 := -\Delta$ in $L^2(\mathbf{R}^d)$, and $H := -\Delta_D$ the Dirichlet Laplacian in Ω with domain $D[H] = H^2(\Omega) \cap H_0^1(\Omega)$. Then for $0 < \sigma < \frac{1}{6}$,

$$\begin{aligned} (\chi_\Omega e^{-tH_0/n} \chi_\Omega)^n(x, y) - e^{-tH}(x, y) &= O(n^{-\sigma}), \\ \text{locally uniformly in } (t, x, y) \in (0, \infty) \times \Omega \times \Omega. \end{aligned} \quad (2.6)$$

We don't know what happens when x or y approaches the boundary of Ω .

Corollary.

$$\| [\chi_\Omega e^{-tH_0/n} \chi_\Omega]^n f - e^{-tH} f \|_{L^2} \rightarrow 0, \quad f \in L^2(\Omega).$$

Consequently, Theorem 2.4 is a *stronger* statement than this corollary, though the latter is also obtained by Kato [29] as an abstract result: If A is a nonnegative

selfadjoint operator and P an orthogonal projection in a Hilbert space $\mathcal{H}$, then $(Pe^{-tA/n}P)^n \rightarrow e^{-tA_P}$, strongly, as $n \rightarrow \infty$, where $A_P := (A^{1/2}P)^*(A^{1/2}P)$. In passing, however, it is an open question whether it holds that $(Pe^{-itA/n}P)^n \rightarrow e^{-itA_P}P$, strongly (Zeno product formula). A partial answer was given in [11].

All Theorems 2.2–2.4 hold with order of products exchanged, e.g. in Theorem 2.2, $[e^{-tV/2n}e^{-tH_0/n}e^{-tV/2n}]^n$ instead of $[e^{-tH_0/2n}e^{-tV/n}e^{-tH_0/2n}]^n$.

Theorem 2.5. (*Ichinose-Tamura 2004* [24])(Unitary Trotter in norm) *Let A and B be selfadjoint, and assume $H := A + B$ to be essentially selfadjoint in a Hilbert space $\mathcal{H}$. Assume that there exists a dense subspace $\mathcal{D}$ of $\mathcal{H}$ with $\mathcal{D} \subset D[A] \cap D[B]$ such that $e^{-itA}, e^{-itB} : \mathcal{D} \rightarrow \mathcal{D}$. Further assume that the commutators $[A, B]$, $[A, [A, B]]$ and $[B, [A, B]]$ are bounded on $\mathcal{H}$. Then*

$$\|(e^{-itB/2n}e^{-itA/n}e^{-itB/2n})^n - e^{-itH}\| = O(n^{-2}), \quad n \rightarrow \infty, \quad (2.7)$$

locally uniformly in the real line $\mathbf{R}$.

As important applications we have ones to the Dirac operator $H = H_0 + V = (i\alpha \cdot \nabla + m\beta) + V(x)$ in $L^2(\mathbf{R}^3)^4$, where $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ and β are the 4 Dirac matrices, with $\partial^\gamma V(x)(|\gamma| = 2)$ being bounded, as well as to the relativistic Schrödinger operator $H = H_0 + V = \sqrt{-\Delta + m^2} + V(x)$ on $L^2(\mathbf{R}^d)$ with $\partial_x^\gamma V(x)$ being bounded for $1 \leq |\gamma| \leq 4$ ($0 \leq |\gamma| \leq 4$, if $m = 0$). In these cases, H are essentially selfadjoint, and satisfy the conditions in the theorem. So it holds that

$$\|[e^{-itV/2n}e^{-itH_0/n}e^{-itV/2n}]^n - e^{-itH}\|_{L^2} = O(n^{-2}), \quad n \rightarrow \infty, \quad (2.8)$$

locally uniformly in $\mathbf{R}$.

However, this theorem does not apply to Schrödinger operators except for the Stark Hamiltonian $(-\Delta + V(x)) + a \cdot x$ in $L^2(\mathbf{R}^d)$, where a is a constant real vector in $\mathbf{R}^d$.

Finally it should be noted that we have shown in Theorems 2.2–2.4 that the convergence is uniform only on compact t -intervals which are away from 0, though in Theorems 2.1 and 2.5, on ones which are allowed to be not away from 0.

3. Optimality of Error Bounds

In this section we discuss optimality of error bounds. The error bound $O(1/n)$ in Theorem 2.1 is optimal, because if both A and B are bounded operators, by the Baker–Campbell–Hausdorff formula we know

$$\begin{aligned} [e^{-tA/n}e^{-tB/n}]^n - e^{-tH} &= R'_n \cdot n^{-1}, \\ [e^{-tB/2n}e^{-tA/n}e^{-tB/2n}]^n - e^{-tH} &= R_n \cdot n^{-2}, \end{aligned}$$

for some R'_n and R_n being uniformly bounded operators which in general are not the zero operator. From this, optimality in the former non-symmetric case is evident. But even in the symmetric case it is optimal. Indeed, there exist unbounded nonnegative selfadjoint operators A, B such that $H = A + B$ is selfadjoint and

$$\|[e^{-tB/2n}e^{-tA/n}e^{-tB/2n}]^n - e^{-tH}\| \geq c(t)n^{-1}$$

for some continuous function $c(t)$ with $c(t) > 0$, $t > 0$ and $c(0) = 0$ ([27]).

However, further in some special symmetric case in Theorem 2.2 where $-\Delta$, V are taken as A , B , we have seen the symmetric product formula hold with a *sharp* error bound $O(n^{-2})$. We can make more precise this result with the 1-dimensional harmonic oscillator $H := H_0 + V := \frac{1}{2}(-\partial_x^2 + x^2)$ in $L^2(\mathbf{R})$.

Theorem 3.1. (*Azuma-Ichinose 2007 [2]*) *There exists bounded continuous functions $C(t) \geq 0$ and $c(t) \geq 0$ in $t \geq 0$, which are positive except $t = 0$ with $C(0) = c(0) = 0$, independent of n , such that for $n = 1, 2, \dots$,*

$$c(t)n^{-2} \leq \| [e^{-\frac{t}{2n}V} e^{-\frac{t}{n}H_0} e^{-\frac{t}{2n}V}]^n - e^{-tH} \| \leq C(t)n^{-2}, \quad t \geq 0. \quad (3.1)$$

This theorem mentions an error bound from below, extending the harmonic oscillator case of Theorem 2.2 which treats only the right-half inequality with $C(t) = C$ being a positive constant depending on each compact t -interval in the open half line $(0, \infty)$.

It is anticipated that the same is true for the Schrödinger operator $H = -\Delta + V(x)$ with growing potentials like $V(x) = |x|^{2m}$ treated in Theorem 2.2.

Theorem 3.1 is obtained as a corollary from the following theorem of its integral kernel version. Here one calculates explicitly the integral kernel $K^{(n)}(t, x, y)$ of $[e^{-tV/2n} e^{-tH_0/n} e^{-tV/2n}]^n$ to estimate its difference from the integral kernel $e^{-tH}(x, y)$ of e^{-tH} .

Theorem 3.2. (*Azuma-Ichinose 2007 [2]*) *There exists a bounded operator $R(t)$ and uniformly bounded operators $\{Q^{(n)}(t)\}_{n=1}^\infty$ with integral kernels $R(t, x, y)$ and $Q^{(n)}(t, x, y)$ being uniformly bounded continuous functions in $(0, \infty) \times \mathbf{R} \times \mathbf{R}$ such that*

$$K^{(n)}(t, x, y) - e^{-tH}(x, y) = [R(t, x, y) + Q^{(n)}(t, x, y)n^{-1}]n^{-2}; \quad (3.2)$$

they satisfy

$$\sup_{x,y} |R(t, x, y)|, \sup_n \sup_{x,y} |Q^{(n)}(t, x, y)| \rightarrow 0, \quad t \rightarrow 0; \quad \sup_{x,y} |R(t, x, y)| \rightarrow 0, \quad t \rightarrow \infty.$$

$R(t, x, y)$ is explicitly given by

$$\begin{aligned} R(t, x, y) = e^{-tH}(x, y) \frac{t^2}{12} & \left[t \left(\frac{1}{4} \frac{e^t + e^{-t}}{e^t - e^{-t}} + \frac{(e^t + e^{-t})xy - (x^2 + y^2)}{(e^t - e^{-t})^2} \right) \right. \\ & \left. + \frac{1}{16} \left(1 + \frac{4xy - (e^t + e^{-t})(x^2 + y^2)}{e^t - e^{-t}} \right) \right]. \quad (3.3) \end{aligned}$$

If $t > 0$, $R(t, x, y)$ can become positive and negative.

Lemma 3.3. $K^{(n)}(t, x, y) =$

$$\begin{aligned}
& \frac{1}{\sqrt{\pi}} \left(\frac{\sqrt{1 + \frac{t^2}{4n^2}}}{\left(1 + \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^n - \left(1 - \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^n} \right)^{1/2} \\
& \times \exp \left[\frac{2\sqrt{1 + \frac{t^2}{4n^2}}}{\left(1 + \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^n - \left(1 - \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^n} xy \right] \\
& \times \exp \left\{ \left[-\frac{t}{4n} - \frac{n}{2t} \left(1 - \frac{\left(1 + \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^{n-1} - \left(1 - \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^{n-1}}{\left(1 + \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^n - \left(1 - \frac{t}{n} \sqrt{1 + \frac{t^2}{4n^2} + \frac{t^2}{2n^2}}\right)^n} \right) \right] (x^2 + y^2) \right\}.
\end{aligned} \tag{3.4}$$

Proof. Calculate the Gaussian integral

$$\begin{aligned}
& K^{(n)}(t, x, y) \\
& \equiv \left(\frac{n}{2\pi t} \right)^{\frac{n}{2}} \int_{\mathbf{R}} \cdots \int_{\mathbf{R}} \prod_{j=1}^n \left[e^{-\frac{t}{4n} x_j^2} e^{-\frac{(x_j - x_{j-1})^2}{2t/n}} e^{-\frac{t}{4n} x_{j-1}^2} \right] dx_1 \cdots dx_{n-1},
\end{aligned}$$

where $x = x_n$, $y = x_0$. We shall encounter with continued fraction to lead to the final expression (3.4) of the lemma.

To show Theorem 3.2, we simply calculate the difference $K^{(n)}(t, x, y) - e^{-tH}(x, y)$, though it is not so simple.

Here we mention what the operator with $R(t, x, y)$ as its integral kernel is. By the Baker–Campbell–Hausdorff formula (e.g. [45], [40]), if A and B are bounded operators, we have

$$\begin{aligned}
& [e^{-tB/2n} e^{-tA/n} e^{-tB/2n}]^n - e^{-t(A+B)} \\
& = \exp \left(-t(A+B) - n^{-2} \frac{t^2}{24} [2A+B, [A, B]] - O_p(n^{-3}) \right) \\
& = e^{-t(A+B)} - n^{-2} \frac{t^2}{24} \int_0^t e^{-(t-s)(A+B)} [2A+B, [A, B]] e^{-s(A+B)} ds + O_p(n^{-3}),
\end{aligned}$$

where $O_p(n^{-3})$ is an operator with norm of $O(n^{-3})$. In our case where $A = -\frac{1}{2}\partial_x^2$, $B = \frac{1}{2}x^2$, we can show $R(t, x, y)$ is just the integral kernel of the operator

$$-\frac{t^2}{24} \int_0^t e^{-(t-s)H} [2H_0 + V, [H_0, V]] e^{-sH} ds,$$

which *does* make sense, though H_0 and V are unbounded operators. We have $[2H_0 + V, [H_0, V]] = -4H_0 + 2V = -4H + 6V$.

4. Idea of Proof

Put $K(\tau) = e^{-\tau B/2} e^{-\tau A} e^{-\tau B/2}$. Note that $0 \leq K(\tau) \leq 1$. Then we need to estimate the difference between $K(t/n)^n$ and e^{-tH} . The general technique of proof is: (i) to establish an appropriate version of Chernoff's theorem ([8]):

$$\begin{aligned} [(1 + \tau^{-1}(1 - K(\tau))^{-1} - (1 + H)^{-1}] &\rightarrow 0, \quad \tau \downarrow 0 \\ \implies [K(t/n)^n - e^{-tH}] &\rightarrow 0, \quad n \rightarrow \infty \end{aligned}$$

and/or (ii) to do telescoping:

$$e^{-tH} - K(t/n)^n = \sum_{k=1}^n e^{-(k-1)tH/n} (e^{-tH/n} - K(t/n)) K(t/n)^{n-k}$$

to estimate each summand on the right. The former method (i) seems to be more efficient than the latter (ii).

In fact, to prove Theorem 1, we use the former method, establishing the following norm version of Chernoff's theorem with error bounds. The case without error bounds was noted by Neidhardt–Zagrebnov [33].

Lemma 4.1. (*Ichinose–Tamura* [23]) *I. Let $\{F(t)\}_{t \geq 0}$ be a family of selfadjoint operators with $0 \leq F(t) \leq 1$, and $H \geq 0$ a selfadjoint operator in a Hilbert space $\mathcal{H}$. Define $S_t := t^{-1}(1 - F(t))$. Then*

(a) *For $0 < \alpha \leq 1$, $\|(1 + S_t)^{-1} - (1 + H)^{-1}\| = O(t^\alpha)$, $t \downarrow 0$ implies*

(b) *For every fixed $\delta > 0$, $\|F(t/n)^n - e^{-tH}\| = \delta^{-2} t^{-1+\alpha} e^{\delta t} O(n^{-\alpha})$, $n \rightarrow \infty$, $t > 0$.*

Therefore for $\alpha = 1$ this convergence is uniform on each compact interval $[0, L]$ in the closed half line $[0, \infty)$.

II. Moreover, in case $H \geq \eta I$ for some constant $\eta > 0$, if for every $\varepsilon > 0$ there exists $\delta(\varepsilon) > 0$ such that $F(t) \leq 1 - \delta(\varepsilon)$ for all $t \geq \varepsilon$, then

$$\|F(t/n)^n - e^{-tH}\| = (1 + 2/\eta)^2 t^{-1+\alpha} O(n^{-\alpha}), \quad n \rightarrow \infty, \quad t > 0.$$

Therefore for $\alpha = 1$ this convergence is uniform on the whole closed half line $[0, \infty)$.

Condition II is satisfied, e.g. for $F(\tau) = e^{-\tau B/2} e^{-\tau A} e^{-\tau B/2}$. For the proof, we refer to [23].

For the proof of Theorems 2.2–2.5 we employ the latter method (ii), and further, for Theorems 2.2–2.4, make a crucial use of Agmon's kernel theorem:

Lemma 4.2. (*Agmon's kernel theorem [1]*)

Let $T : L^2(\mathbf{R}^d) \rightarrow L^2(\mathbf{R}^d)$ be a bounded operator with ranges of T and its adjoint T^* satisfying $R[T], R[T^*] \subset H^m(\mathbf{R}^d)$, $m > d$. If T is an integral operator with integral kernel $T(x, y)$ being a bounded continuous function in $\mathbf{R}^d \times \mathbf{R}^d$ such that

$$(Tf)(x) = \int T(x, y)f(y) dy, \quad f \in L^2,$$

then

$$|T(x, y)| \leq C(\|T\|_m + \|T^*\|_m)^{\frac{d}{m}} \|T\|^{1-\frac{d}{m}},$$

where $\|T\|_m := \|T\|_{\mathcal{L}(L^2 \rightarrow H^m)}$ is the operator norm of T as a bounded operator of $L^2(\mathbf{R}^d)$ into the Sobolev space $H^m(\mathbf{R}^d)$.

Indeed, we estimate the $\mathcal{L}(L^2 \rightarrow H^m)$ -operator norm of the difference $T = [e^{-tV/2n} e^{t\Delta/n} e^{-tV/2n}]^n - e^{t(-\Delta+V)}$.

5. Concluding Remarks

We have so far considered the case where the operator sum $H = A + B$ of two nonnegative selfadjoint operators A and B is selfadjoint. However, otherwise, the exponential product formula in norm does not in general hold for the form sum $H = A + B$ of two selfadjoint operators $A \geq 0$, $B \geq 0$, even if it is essentially selfadjoint on $D[A] \cap D[B]$ (see [43]). Nevertheless, there is some case where it holds:

Theorem 5.1. (*Ichinose-Neidhardt-Zagrebnov 2004 [16]*) Let $H = A + B$ be the form sum of A and B . If $D[H^\alpha] \subseteq D[A^\alpha] \cap D[B^\alpha]$ for some $\frac{1}{2} < \alpha < 1$, and $D[A^{\frac{1}{2}}] \subseteq D[B^{\frac{1}{2}}]$, then

$$\| [e^{-tB/2n} e^{-tA/n} e^{-tB/2n}]^n - e^{-tH} \| = O(n^{-(2\alpha-1)}), \quad (5.1)$$

$$\| [e^{-tA/n} e^{-tB/n}]^n - e^{-tH} \| = O(n^{-(2\alpha-1)}), \quad (5.2)$$

locally uniformly in $[0, \infty)$.

This error bound in (5.1)/(5.2) is also optimal. For this we refer to [43]. The condition for the domains of A and B is not symmetric. It is an open question whether one may improve it so as to become symmetric with respect to A and B .

Finally, as we should like to mention, there is a very nice Feynman path integral formula which represents the Schrödinger semigroup, called the *Feynman-Kac formula*

$$\begin{aligned} (e^{-tH} f)(x) &= (e^{-t(-\Delta+V)} f)(x) \\ &= \int_{B \in C([0, \infty) \rightarrow \mathbf{R}^d), B(0)=x} \exp[-\int_0^t V(B(s)) ds] f(B(t)) d\mu(B), \end{aligned}$$

where $\mu(\cdot)$ is the Wiener measure on the path space $C([0, \infty) \rightarrow \mathbf{R}^d)$ (e.g. ([39])). We may use this formula to get whatever results, in fact, a lot of them. This is

a big advantage! But disadvantage is that it is only restricted to the Schrödinger operator or Laplacian. For instance, if we think of the semigroup for the relativistic Schrödinger operator $H = \sqrt{-\Delta + m^2} + V(x)$, we have to establish another Feynman–Kac formula (cf. [28]).

Indeed, the Feynman–Kac formula is one of the realizations of Feynman path integral as a *true integral* on a path space. However, as Nelson [36] noted, the exponential/Trotter product formula also can give a meaning to the Feynman path integral as a *time-sliced approximation* by finite-dimensional integrals (cf. [15]). What it has advantage at is that we may apply it to the sum $H = A + B$ of any two selfadjoint operators A, B bounded from below.

Acknowledgements. One of the authors (T.I.) would like to thank Professor Vadim Adamyan for kind and warm hospitality during his stay in Odessa for the Mark Krein Centenary Conference, April 2007. This work (of T.I. and H.T.) was partially supported by Grant-in-Aid for Exploratory Research No. 17654030 and by Grant-in-Aid for Scientific Research (B) No. 18340049, Japan Society for the Pomotion of Sciences.

References

- [1] S. Agmon, *On kernels, eigenvalues, and eigenfunctions of operators related to elliptic problems*, Comm. Pure Appl. Math. **18** (1965), 627–663.
- [2] Y. Azuma and T. Ichinose, *Note on norm and pointwise convergence of exponential products and their integral kernels for the harmonic oscillator*, to appear in Integral Equations Operator Theory (published on line first Nov. 12, 2007).
- [3] V. Cachia, H. Neidhardt and V. A. Zagrebnov, *Accretive perturbations and error estimates for the Trotter product formula*, Integral Equations Operator Theory **39** (2001), 396–412.
- [4] V. Cachia, H. Neidhardt and V. A. Zagrebnov, *Comments on the Trotter product formula error-bound estimates for nonself-adjoint semigroups*, Integral Equations Operator Theory **42** (2002), 425–448.
- [5] V. Cachia and V. A. Zagrebnov, *Operator-norm convergence of the Trotter product formula for sectorial generators*, Lett. Math. Phys. **50** (1999), 203–211.
- [6] V. Cachia and V. A. Zagrebnov, *Trotter product formula for nonself-adjoint Gibbs semigroups*, J. London Math. Soc. (2) **64** (2001), 436–444.
- [7] V. Cachia and V. A. Zagrebnov, *Operator-norm convergence of the Trotter product formula for holomorphic semigroups*, J. Operator Theory **46** (2001), 199–213.
- [8] P.R. Chernoff, *Product Formulas, Nonlinear Semigroups, and Addition of Unbounded Operators*, Memoirs Amer. Math. Soc., **140**, 1974.
- [9] B. O. Dia and M. Schatzman, *An estimate on the Kac transfer operator*, J. Functional Analysis **145** (1997), 108–135.
- [10] A. Doumeki, T. Ichinose and H. Tamura, *Error bounds on exponential product formulas for Schrödinger operators*, J. Math. Soc. Japan **50** (1998), 359–377.

- [11] P. Exner and T. Ichinose, *A product formula related to quantum Zeno dynamics*, Ann. H. Poincaré **6** (2005), 195–215.
- [12] I.C. Gohberg and M.G. Krein, *Introduction to the theory of linear non-selfadjoint operators (in Hilbert space)*, Translations of Mathematical Monographs, Vol. 18, Amer. Math. Soc., Providence, R.I. 1969; its Russian edition, Izdat. “Nauka”, Moscow 1965.
- [13] B. Helffer, *Around the transfer operator and the Trotter–Kato formula*, Operator Theory: Advances and Appl. **79** (1995), 161–174.
- [14] F. Hiai, *Trace norm convergence of exponential product formula*, Lett. Math. Phys. **33** (1995), 147–158.
- [15] T. Ichinose, *Time-sliced approximation to path integral and Lie–Trotter–Kato product formula*, In: “A Garden of Quanta”, Essays in Honor of Hiroshi Ezawa for his seventieth birthday, World Scientific 2003, pp. 77–93.
- [16] T. Ichinose, H. Neidhardt and V. A. Zagrebnov, *Trotter–Kato product formula and fractional powers of self-adjoint generators*, J. Functional Analysis **207** (2004), 33–57; *Operator norm convergence of Trotter–Kato product formula*, Ukrainian Mathematics Congress—2001 (Ukrainian), 100–106, Natsional. Akad. Nauk Ukraini, Inst. Mat., Kiev, 2002.
- [17] T. Ichinose and S. Takanobu, *Estimate of the difference between the Kac operator and the Schrödinger semigroup*, Commun. Math. Phys. **186** (1997), 167–197.
- [18] T. Ichinose and S. Takanobu, *The norm estimate of the difference between the Kac operator and the Schrödinger semigroup: A unified approach to the nonrelativistic and relativistic cases*, Nagoya Math. J. **149** (1998), 51–81.
- [19] T. Ichinose and S. Takanobu, *The norm estimate of the difference between the Kac operator and the Schrödinger semigroup II: The general case including the relativistic case*, Electronic Journal of Probability **5** (2000), Paper 5, pages 1–47; URL <http://www.math.washington.edu/~ejpecp/>
- [20] T. Ichinose and H. Tamura, *Error estimate in operator norm for Trotter–Kato product formula*, Integral Equations Operator Theory **27** (1997), 195–207.
- [21] T. Ichinose and H. Tamura, *Error bound in trace norm for Trotter–Kato product formula of Gibbs semigroups*, Asymptotic Analysis **17** (1998), 239–266.
- [22] T. Ichinose and H. Tamura, *Error estimate in operator norm of exponential product formulas for propagators of parabolic evolution equations*, Osaka J. Math. **35** (1998), 751–770.
- [23] T. Ichinose and H. Tamura, *The norm convergence of the Trotter–Kato product formula with error bound*, Commun. Math. Phys. **217** (2001), 489–502; Erratum, *ibid.* **254** (2005), 255.
- [24] T. Ichinose and H. Tamura, *Note on the norm convergence of the unitary Trotter product formula*, Lett. Math. Phys. **70** (2004), 65–81.
- [25] T. Ichinose and H. Tamura, *Sharp error bound on norm convergence of exponential product formula and approximation to kernels of Schrödinger semigroups*, Comm. Partial Differential Equations **29** (2004), 1905–1918.
- [26] T. Ichinose and H. Tamura, *Exponential product approximation to integral kernel of Schrödinger semigroup and to heat kernel of Dirichlet Laplacian*, J. Reine Angew. Math. **592** (2006), 157–188.

- [27] T. Ichinose, H. Tamura, Hiroshi Tamura and V. A. Zagrebnov, *Note on the paper “The norm convergence of the Trotter–Kato product formula with error bound” by Ichinose and Tamura*, Commun. Math. Phys. **221** (2001), 499–510.
- [28] T. Ichinose and Hiroshi Tamura, *Imaginary-time path integral for a relativistic spinless particle in an electromagnetic field*, Commun. Math. Phys. **105**, 239–257(1986); For the final result of this see: T. Ichinose, *Some results on the relativistic Hamiltonian: Selfadjointness and imaginary-time path integral*, Differential Equations and Mathematical Physics, pp. 102–116, International Press, Boston 1995.
- [29] T. Kato, *Trotter’s product formula for an arbitrary pair of self-adjoint contraction semigroups*, In: Topics in Functional Analysis (essays dedicated to M. G. Krein on the occasion of his 70th birthday), edited by I. Gohberg and M. Kac, Academic Press, New York 1978, pp.185–195.
- [30] H. Neidhardt and V. A. Zagrebnov, *The Trotter–Kato product formula for Gibbs semigroups*, Commun. Math. Phys. **131** (1990), 333–346.
- [31] H. Neidhardt and V. A. Zagrebnov, *On the Trotter product formula for Gibbs semigroups*, Ann. Physik (7)**47** (1990), 183–191.
- [32] H. Neidhardt and V. A. Zagrebnov, *On error estimates for the Trotter–Kato product formula*, Lett. Math. Phys. **44** (1998), 169–186.
- [33] H. Neidhardt and V. A. Zagrebnov, *Trotter–Kato product formula and operator-norm convergence*, Commun. Math. Phys. **205** (1999), 129–159.
- [34] H. Neidhardt and V. A. Zagrebnov, *Trotter–Kato product formula and symmetrically normed ideals*, J. Functional Analysis **167** (1999), 113–147.
- [35] H. Neidhardt and V. A. Zagrebnov, *Fractional powers of self-adjoint operators and Trotter–Kato product formula*, Integral Equations Operator Theory **35** (1999), 209–231.
- [36] E. Nelson, *Feynman integrals and the Schrödinger equation*, J. Math. Phys. **5** (1964), 332–343.
- [37] Dzh. L. Rogava, *Error bounds for Trotter–type formulas for self-adjoint operators*, Functional Analysis and Its Applications **27** (1993), 217–219.
- [38] R. Schatten, Norm ideals of completely continuous operators, Springer, Berlin 1960.
- [39] B. Simon, Functional Integration and Quantum Physics, Academic Press, London 1979.
- [40] M. Suzuki, *On the convergence of exponential operators — the Zassenhaus formula, BCH formula and systematic approximations*, Commun. Math. Phys. **57** (1977), 193–200.
- [41] S. Takanobu, *On the estimate of the integral kernel for the Trotter product formula for Schrödinger operators*, Ann. Probab. **25** (1997), 1895–1952.
- [42] S. Takanobu, *On the trace norm estimate of the Trotter product formula for Schrödinger operators*, Osaka J. Math. **35** (1998), 659–682.
- [43] Hiroshi Tamura, *A remark on operator-norm convergence of Trotter–Kato product formula*, Integral Equations Operator Theory **37** (2000), 350–356.
- [44] H. F. Trotter, *On the product of semigroups of operators*, Proc. Amer. Math. Soc. **10** (1959), 545–551.

- [45] S. Varadarajan, *Lie Groups, Lie Algebras, and Their Representations*, Springer, Berlin–Heidelberg–New York–Tokyo 1974, 1984.
- [46] V. A. Zagrebnov, *The Trotter–Lie product formula for Gibbs semigroups*, J. Math. Phys. **29** (1988), 888–891.
- [47] V. A. Zagrebnov, *Topics in the Theory of Gibbs Semigroups*, Leuven Notes in Mathematical and Theoretical Physics, Vol. 10, Leuven University Press 2003.

Takashi Ichinose

Department of Mathematics, Kanazawa University, Kanazawa, 920–1192, Japan

e-mail: ichinose@kenroku.kanazawa-u.ac.jp

Hideo Tamura

Department of Mathematics, Okayama University, Okayama, 700–8530, Japan

e-mail: tamura@math.okayama-u.ac.jp